

Phase II clinical trial in DMD Fully Enrolled

- Enrolment completed with the last patient into the trial to be dosed this month
- No Serious Adverse Events reported to date
- Dosing to be completed early November 2019

Antisense Therapeutics ("ANP" or the "Company") is pleased to advise that the Phase II clinical trial of its immunomodulatory therapy, ATL1102 for Duchenne Muscular Dystrophy (DMD) is now fully enrolled.

The open label six-month dosing trial of ATL1102 in nine non-ambulant patients with DMD aged between 10 and 18 years is being conducted at the neuromuscular centre of the Royal Children's Hospital in Melbourne which operates the largest clinic in the southern hemisphere treating children with DMD.

Three patients have completed their 24 weeks of dosing. Four patients are in the treatment phase of the study. The final two patients have been screened and are scheduled to commence their dosing before the end of this month. No Serious Adverse Events have been reported to date. Dosing of all patients in the trial is to be completed in early November 2019.

ATL1102 is being developed as a novel treatment for the inflammation that exacerbates muscle fibre damage in DMD patients, currently treated with corticosteroids. Corticosteroids have a range of serious side effects when used for a prolonged period as required in DMD. Currently there are a limited number of anti-inflammatory drugs in clinical development for DMD.

The primary endpoints of the trial relate to the safety and tolerability of ATL1102. The efficacy of ATL1102 will also be assessed in terms of its effects on disease processes and progression. The extended dosing period (24 weeks) of this clinical trial may allow for ATL1102 to show an improvement in the clinical endpoints that are relevant to DMD disease progression (e.g. the upper limb function of the boys) and that are of the type that would be required for future product registration. Previous published studies in non-ambulant DMD boys have reported that after six months there is a significant decline in such functional endpoints even if the patients are on corticosteroids. One of the objectives of the ATL1102 trial is to address the decline in upper limb strength and function of the non-ambulant DMD patients.

The Company expects to report trial results shortly after the completion of dosing. As previously advised, the Phase II DMD clinical trial is an open label study and there may be an opportunity for non-statistical study read-outs on preliminary data prior to the completion of dosing in all patients. This would require a sufficient number of patients to have completed their dosing for the Company to be confident and certain of the robustness of such results for disclosure.

Mark Diamond, CEO of Antisense Therapeutics said: " We are very pleased to have achieved the key clinical trial milestone of completion of patient enrolment into our Phase II clinical trial of ATL1102 in DMD patients. The intensive safety monitoring required for this first study in boys with DMD requires frequent visitations to the hospital over the course of the trial period which is particularly challenging for wheelchair bound boys and their families and so we greatly appreciate their commitment to participate in this trial. The Data Safety Monitoring Board have been periodically evaluating the safety related trial data and have on each occasion recommended continuation of the trial with no safety concerns. We are now set to have dosing completed and trial results reported in the fourth quarter of this year, which presents as a very exciting prospect for the Company and potentially for future treatment of DMD patients".

For more information please contact:

Antisense Therapeutics

Mark Diamond
Managing Director
+61 (0)3 9827 8999
www.antisense.com.au

Investment Enquiries

Gennadi Koutchin
XEC Partners
gkoutchin@xecpartners.com.au
1300 932 037

About Antisense Therapeutics Limited (ASX: ANP) is an Australian publicly listed biopharmaceutical company, developing and commercialising antisense pharmaceuticals for large unmet markets. The products are in-licensed from Ionis Pharmaceuticals Inc. (NASDAQ:IONS), world leaders in antisense drug development and commercialisation. ATL1102 (injection) has successfully completed a Phase II efficacy and safety trial, significantly reducing the number of brain lesions in patients with relapsing-remitting multiple sclerosis (RRMS). ATL1103 drug designed to block GHr production successfully reduced blood IGF-I levels in Phase II clinical trials in patients with the growth disorder acromegaly. The Company is conducting a Phase II clinical trial of ATL1102 in DMD patients at the Royal Childrens Hospital, Melbourne.

About ATL1102 ATL1102 is an antisense inhibitor of CD49d, a subunit of VLA-4 (Very Late Antigen-4). Antisense inhibition of VLA-4 expression has demonstrated activity in a number of animal models of inflammatory disease including asthma and MS with the MS animal data having been published in a peer reviewed scientific journal. ATL1102 was shown to be highly effective in reducing MS lesions in a Phase IIa clinical trial in RR-MS patients. The ATL1102 Phase IIa clinical data has been published in the medical Journal *Neurology* (Limmroth, V. et al Neurology, 2014; 83(20): 1780-1788).

About DMD Duchenne Muscular Dystrophy (DMD) is an X-linked disease that affects 1 in 3600 to 6000 live male births (Bushby *et al*, 2010). DMD occurs as a result of mutations in the dystrophin gene which causes a substantial reduction in or absence of the dystrophin protein. Children with DMD have dystrophin deficient muscles and are susceptible to contraction induced injury to muscle that triggers the immune system which exacerbates muscle damage as summarized in a publication co-authored by the Director of the FDA CDER (Rosenberg et al, 2015). Ongoing deterioration in muscle strength affects lower limbs leading to impaired mobility, and also affects upper limbs, leading to further loss of function and self-care ability. The need for wheelchair use can occur in early teenage years for patients on corticosteroids with a mean age of 13, with respiratory, cardiac, cognitive dysfunction also emerging. Patients with a greater number of immune T cells expressing high levels of CD49d have more severe and progressive disease and are wheelchair bound by the age of 10 despite being on corticosteroid treatment (Pinto Mariz et al, 2015). With no intervention, the mean age of life is approximately 19 years. The management of the inflammation associated with DMD is currently addressed via the use of corticosteroids, however they are acknowledged as providing insufficient efficacy and are associated with significant side effects. As a consequence, there is an acknowledged high need for new therapeutic approaches for the treatment of inflammation associated with DMD.

Rosenberg AS, Puig M, Nagaraju K, *et al*. Immune-mediated pathology in Duchenne muscular dystrophy. *Sci Transl Med* 2015, 7: 299rv4.

Busby et al for the DMD Care Consideration Working Group/ *Diagnosis and management of Duchenne muscular dystrophy, part 1* Lancet Neurol. **2010** Jan;9(1):77-93 and *part 2* Lancet Neurol. **2010** Feb;9(2):177-89 .

*Pinto-Mariz F, Carvalho LR, Araújo AQC, *et al*. CD49d is a disease progression biomarker and a potential target for immunotherapy in Duchenne muscular dystrophy. *Skeletal Muscle* 2015, 5: 45-55